

PARTIAL ALBINOS AND WHITE BLAZE INDIVIDUALS OF SOUTHERN RIGHT WHALES (*Eubalaena australis*) IN BRAZILIAN WATERS.

Paulo A.C. Flores<sup>1,2</sup>; Karina R. Groch<sup>1</sup>; Paulo H. Ott<sup>3,4</sup> and Márcia Engel<sup>5,6</sup>

Affiliations and addresses:

<sup>1</sup> International Wildlife Coalition Brazil, P.O. Box 5087, Florianópolis, SC, Brasil 88040-970

<sup>2</sup> Laboratório de Dinâmica Populacional, Pós Graduação em Biociências, PUCRS, Av. Ipiranga 6681, Porto Alegre, RS, Brasil 90619-900

<sup>3</sup> GEMARS and CECLIMAR/UFRGS, R. Felipe Neri, 382/203, Porto Alegre, RS 90440-150, Brasil

<sup>4</sup> PPG Genética e Biologia Molecular, Depto de Genética, UFRGS, P.O. Box 15053, Porto Alegre, RS, Brasil 91501-970

<sup>5</sup> Projeto Baleia Jubarte/Instituto Baleia Jubarte, Praia do Kitongo s/n, Caravelas, BA, Brasil 45900-000

<sup>6</sup> Pós-Graduação em Biocências, PUCRS, Av. Ipiranga 6681, Porto Alegre, RS, Brasil 90619-900

ABSTRACT

Partial albinism (or gray morphs) in southern right whales (*Eubalaena australis* Desmoulins 1822) is not rare in individuals of Argentinean, South African and Western Australian populations. Right whales in Brazilian waters are distributed from southern Rio Grande do Sul state (~ 34°S) to the Arolhos Bank, Bahia state (~ 18° S), during the species breeding and nursing period from June to November. Right whales are common in Rio Grande do Sul and Santa Catarina, the main concentration area for the Brazilian population, while more rare in the northern areas. Despite research effort has been directed to the species along the southern Brazilian coast since 1982, including photo-identification studies from 1986 onwards, just recently have the presence of partial albinism come to attention in this area. The first register was of a live calf photographed with its mother in Arolhos Bank (17°55'S, 38°56'W) in October 1999. A second, distinct calf was sighted in August 2000 off Santa Catarina (28°28'S) and then followed and photographed during 45 days, providing information on behavior and movement patterns. The only previous record of an albino right whale along the Brazilian coast was based on a photograph of a dead calf stranded off Rio Grande do Sul in 1982. White blaze animals were also recorded during ongoing studies along Santa Catarina coast: one calf in 1987, one calf in 1999 and two adults in 2000. The actual frequency of partial albinism, white blaze individuals and animals with white ventral patches should be evaluated through the analysis of the photographic catalogue in the Brazilian population and compared with the data from others breeding/calving grounds. These analyses in conjunction with underway genetic studies could help to understand the relationship among the different South Atlantic right whale populations.

Anomalously white cetaceans have been recorded in 20 species (Hain and Leatherwood 1982, Fertl *et al.* 1999), seeming relatively common in the southern right whale, *Eubalaena australis*, populations (Schaeff *et al.* 1999). Dorsal skin color patterns in *E. australis* also include white-blaze, gray-blaze, gray-and-white-blaze and the typical black (Best 1990, Payne *et al.* 1983).

Although *E. australis* is common in southern Brazil (especially Santa Catarina coast) with some occurring in southeastern and very few on northeastern (around Arolhos Bank) (Fig. 1), no records or comments on the occurrence of partial albinistic animals have been made (Pinedo and Castello 1979, Lodi *et al.* 1996, Simões-Lopes *et al.* 1992, IWC/Brazil 1999, Groch 2000, Greig *et al.* 2001, Santos *et al.* 2001). Here we present information on partial albinistic animals as well as gray-blaze and white-blaze individuals in the coast of Brazil.

Three partial albinos have been recorded (two in southern Brazil, Figs. 2A, B and D, and another in northeastern, Fig. 2C), covering roughly the whole distribution of the species in Brazil (Fig. 1). A higher number of white-blaze (Fig. 2E) and gray-blaze (Figs. 2B and F) were observed but only in Santa Catarina, where photoidentification from aerial surveys, land based observations and boat surveys

including biopsy efforts have been conducted since 1987. This season of 2001 another partial albino, two white-blaze and two gray-blaze individuals were seen in Santa Catarina.

While for two partial albino calves no information is available, one partial albino calf was born from a gray-morph mother (Fig. 2C). The total number and proportion of individuals with dorsal skin color patterns is still under analysis (KRG Ph.D. thesis), together with genetic evaluation might help us to enlighten the relationship of Brazilian right whales to other populations in the Southern Hemisphere.

#### Acknowledgments

Field work and laboratory were funded by World Wildlife Fund – US, Whale and Dolphin Conservation Society, The Barbara Delano Foundation, Island Foundation, Schlösser Textile Co., World Wildlife Fund – Brazil, and the International Wildlife Coalition in Sanata Catarina; Petrobrás and IBAMA in Abrolhos region, Bahia. CAPES and CNPq/National Graduation Programs also gave partial funds through scholarships to PACF (CNPq proc. n° 146609/1999-9), KRG (CAPES) and PHO (CNPq proc. n° 144064/98-7). Logistical support provided by Turismo Vida, Sol & Mar (Imbituba, SC), local offices of IBAMA in Florianópolis and Laguna, SC as well as in Torres, RS.

#### References

- Best, P.B., Payne, R., Rowntree, V., Palazzo, J.T. and Both, M.C. 1993. Long range movements of South Atlantic right whales *Eubalaena australis*. *Mar. Mamm. Sci.* 9: 227-234.
- Fertl, D., Pusser, L.T. and Long, J.J. 1999. First record of an albino bottlenose dolphin (*Tursiops truncatus*) in the Gulf of Mexico, with a review of anomalously white cetaceans. *Mar. Mamm. Sci.* 15(1): 227-234.
- Greig, A.B., Secchi, E.R., Zerbini, A.N. and Dalla-Rosa, L. 2001. Stranding events of southern right whales, *Eubalaena australis*, in southern Brazil. *J. Ceacean. Res. Manage.* (Special Issue) 2: 157-160.
- Groch, K.R. 2000. Ocupação preferencial de áreas de concentração pela Baleia Franca Austral, *Eubalaena australis* (Desmoulins, 1822), CETACEA, MYSTICETI, no litoral sul do Brasil. Master Thesis, Universidade Federal do Rio Grande do Sul. 61pp. + appendices. [In Portuguese]
- Hain, H.W. and Leatherwood, S. 1982. Two sightings of white pilot whales, *Globicephala melaena*, and summarize records of anomalously white cetaceans. *J. Mamm.* 63: 338-343.
- International Wildlife Coalition/Brazil. 1999. Plano de Ação para a Conservação da Baleia Franca, *Eubalaena australis*, em Santa Catarina, Brasil. International Wildlife Coalition/Brazil. 59pp. [In Portuguese]
- Lodi, L., Siciliano, S. and Bellini, C. 1996. Ocorrências e conservação de baleias francas do sul no litoral do Brasil. *Papéis Avulsos de Zool., São Paulo*, 39(17): 307-328.
- Payne, R., Brazier, O., Dorsey, E.M., Perkins, J.S., Rowntree, V.J. and Titus, A. (1983). External features in southern right whales (*Eubalaena australis*) and their use in identifying individuals: 371-445 In *Communication and behavior of whales*. R. Payne (ed.): Westview Press, Boulder, CO.
- Pinedo, M.C. and Castello, H.P. 1979. Southern right whales (*Eubalaena australis*) along the Southern Brazilian coast. *J. Mammalogy* 60(2): 429-430.
- Santos, M.C. O., Siciliano, S., Souza, S.P. and Pizzorno, J.L.A. 2001. Occurrence of southern right whales (*Eubalaena australis*) along southeastern Brazil. *J. Ceacean. Res. Manage.* (Special Issue) 2: 153-156.
- Schaeff, C.M., Best, P.B., Rowntree, V.J., Payne, R., Jarvis, C. and Portway, V.A. 1999. Dorsal skin color patterns among southern right whales: (*Eubalaena australis*): Genetic basis and evolutionary significance. *The J. Heredity* 90(4): 464-471.
- Simões-Lopes, P.C., Palazzo Jr., J.T., Both, M.C. and Ximenez, A. 1992. Identificação, movimentos e aspectos biológicos da baleia franca austral (*Eubalaena australis*) na costa sul do Brasil. Proceedings of the 3<sup>o</sup> Reunión de Trabajo de Especialistas en Mamíferos Acuáticos de América del Sur, Montevideo, 1988. 62-66pp. [Available from palazzo@zaz.com.br]